

Catalogue No.

AB0291-100

Qty:

100 µg

Anti-Nucleolin

Source: Goat

General description: Goat polyclonal antibody to Nucleolin. NCL is a eukaryotic nucleolar phosphoprotein involved in the synthesis and maturation of ribosomes. It is found associated with pre-ribosomal particles and intranucleolar chromatin. May play a role in the process of transcriptional elongation and chromatin decondensation by binding to histone H1.

Alternative names: B530004O11Rik, C23, D0Nds28, D1Nds28, Nucl antibody.

Form: Polyclonal antibody supplied as a 100 µl (1 mg/ml) aliquot in PBS, 20% glycerol and 0.05% sodium azide. This antibody is epitope-affinity purified from goat antiserum.

Immunogen: Purified recombinant peptide derived from within residues 500 aa to the C-terminus of human NCL produced in E. coli.

Specificity: Detects endogenous levels of NCL by Western blot.

Reactivity: Reacts with Human, Rat, Mouse, Monkey and Canine proteins

Sample	WB	IHC (F)	IHC (P)	IF	ELISA
Human	+++	ND	ND	ND	ND
Rat	+++	ND	ND	ND	ND
Mouse	+++	ND	ND	ND	ND
Canine	+++	ND	ND	ND	ND
Monkey	+++	ND	ND	ND	ND


+++ excellent, ++ good, + poor, ND not determined

Usage:

WB: 1:1,000-1:5,000

Storage: For continuous use, store at 2-8 C for one-two days. For extended storage, store in -20 C freezer. Working dilution samples should be discarded if not used within 12 hours.

Special instructions: The antibody solution should be gently mixed before use..



Anti-NCL Ab at 1/2,500 dilution; lysates at 50 µg per lane; rabbit polyclonal to goat IgG (HRP) at 1/10,000 dilution;

For research use only, not for diagnostic use

SICGEN's Proprietary Immunogen Policy

In order to produce high specific antibodies SICGEN has invested a lot of time and effort into selecting immunogen sequences. SICGEN has decided to protect this information by not publishing it on the website. However, these sequences are available on request.