SICGEN ANTIBODIES

Product Data Sheet

Catalogue No.	Qty:
AB0067-200	400 µg

Anti-GAPDH

Source: Goat

General description: Goat polyclonal to GAPDH (glyceraldehyde 3-phosphate dehydrogenase). GAPDH catalyzes an important energy-yielding step in carbohydrate metabolism, the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD). The enzyme exists as a tetramer of identical chains.

Alternative names: glyceraldehyde 3-phosphate dehydrogenase, glyceraldehyde-3-phosphate dehydrogenase, G3PD, GAPD, HGNC:4141, GAPDH antibody.

Form: Polyclonal antibody supplied as a 200 µl (2 mg/ml) aliquot in PBS, 20% glycerol and 0.05% sodium azide. This antibody is epitope-affinity purified from goat antiserum.

Immunogen: Purified recombinant peptide derived from within residues 240 aa to the C-terminus of human GAPDH produced in E. coli.

Specificity: Detects a band of 37 kDa by Western blot in the following human (293A, HMEC-1, U-118, HaCat), rat (TR-iBRB), mouse (AtT-20, Hepa), canine (D17) and monkey (COS-7) whole cell lysates.

Reactivity: Reacts with Human, Rat, Mouse, Monkey and Canine proteins

Sample	WB	IHC (F)	IHC (P)	IF	ELISA
Human	+++	+++	+++	+++	ND
Rat		+++	+++	+++	ND
Mouse	+++	+++	+++	+++	ND
Monkey	+++	+++	+++	+++	ND
Canine	+++	+++	+++	+++	ND

+++ excellent, ++ good, + poor, ND not determined

Usage:

WB: 1:500-1:10,000


IF:	1:50-1:250
IHC (P):	1:200-1:1,000
IHC (F):	1:200-1:1,000

Storage: For continuous use, store at 2-8 C for one-two days. For extended storage, store in -20 C freezer. Working dilution samples should be discarded if not used within 12 hours.


Special instructions: The antibody solution should be gently mixed before use..

References:

1. Ramos S, Ademolue TW, Jentho E, et al. Cell Metab 2022 Aug PMID: 35841892
2. Broughton K, Korski K, Echeagaray O, et al. Gene Ther. 2019 Aug PMID: 31239537
3. Korski KI, Kubli DA, Wang BJ, et al. Stem Cells. 2019 Jan PMID: 30629785
4. Broughton K, Korski K, Echeagaray O, et al. bioRxiv 521732
5. Korski KI,MSc Thesis, Univ San Diego, USA, 2018
6. Choudhury SS, MSc Thesis, Univ San Diego, USA, 2018
7. Sacchi V, Wang BJ, Kubli D, et al. J Am Heart Assoc. 2017 Oct (Supplemental Information) PMID: 29018025
8. Khalafalla FG, Kayani W, Kassab A, et al. J Physiol. 2017 Oct PMID: 28980705
9. Firouzi F, MSc Thesis, San Diego State University, USA, 2017
10. Gonçalves A, Lin CM, Muthusamy A, et al. Invest Ophthalmol Vis Sci. 2016 May PMID: 27163772
11. Hyenne V, Apaydin A, Rodriguez D, et al. J Cell Biol. 2015 Oct PMID: 26459596
12. Helsby MA, Leader PM, Fenn JR. BMC Cell Biol. 2014 Feb PMID: 24528853
13. Grilo MLJ, MSc Thesis, University of Coimbra, Portugal, 2014
14. Matos CTSM, PhD Thesis, NOVA University of Lisbon, Portugal, 2013



Anti-GADPH Ab at 1/1,000 dilution; lysates at 100 µg per lane; rabbit polyclonal to goat IgG (HRP) at 1/10,000 dilution;



Immunofluorescence – anti-GAPDH Ab in Hepa1-6 cells at 1/50 dilution; cells were fixed with methanol;

For research use only, not for diagnostic use

SICGEN's Proprietary Immunogen Policy

In order to produce high specific antibodies SICGEN has invested a lot of time and effort into selecting immunogen sequences. SICGEN has decided to protect this information by not publishing it on the website. However, these sequences are available on request.